



READ ALL OF THIS INFORMATION LEAFLET CAREFULLY BEFORE YOU START USING THIS

PRODUCT

This product is intended only for adult smokers. This product is not recommended for non-smokers, young people (under the age of 18), pregnant or breast-feeding women and those experiencing general ill health.

INTRODUCTION

- Nicotine Salts is a specific formulation of nicotine used in low wattage devices.
 - Nicotine Salts offer high nicotine strength and are available in 10mg and 20mg.
 - Nicotine Salts are made to have a smoother throat hit even at high
- nicotine strength.

 Beginner users need to start using at low nicotine level

WARNING AND PRECAUTIONS

Nicotine containing e-liquids are not recommended for:

recommended

vears

- Non-smokers
 Young people under the age of 18
- Pregnant or breast feeding women, as nicotine is known to cross the placenta and to be excreted breast milk
- Those experiencing general ill health, such as heart disease, stomach and duodenal ulcers, liver or kidney problems, long term throat disease or difficulty breathing due to bronchitis. emphyseme or asthma
- Those who have an overactive thyroid gland or have a phaeochromocytoma (a tumour of the adrenal gland that can affect high blood pressure).
- Those who have diabetes as nicotine may affect your blood sugar levels

Those taking the following medication:

 Theophylline (to treat breathing problems)

Ropinirole (to treat restless legs)
 Clozapone (to treat schizophrenia)

INDICATIONS FOR USE

Shake well before use



Push the cap down and turn it anti-clockwise at the same time. Remove the cap.



Insert the nozzle of the bottle into the filling port of the cartridge. The nozzle width is 1.8 to 2.6mm.



Squeeze the bottle gently to fill the cartridge.



Carefully pull the nozzle out and put the cap back. Store the product in a safe and dry place.

POSSIBLE SIDE EFFECTS

If you inhale too much nicotine or use the product too much you may experience one or more of the following effects:

- Feeling faint
- Nausea
- Headache
- Coughing
- Irritation in the mouth or throat
- Dizziness
 Stomach discomfort
- Hiccups - Nasal congestion
- Vomiting (being sick)
 Chest palpitations
- Abnormal heart rate

If you experience any of these side effects not listed in this leaflet, discontinue using this product. If the effect(s) continue after you stop use, consult your doctor or healthcare professional and bring this leaflet to their attention.

ADDICITIVENESS

Nicotine is highly addictive and is therefore intended for adult smokers only.

TOXICITY

This product contains nicotine which may produce toxic effects if consumed orally, especially in infants and children. Please DO NOT drink this product. In case it is consumed, please take this leaflet with you, and see a general practitioner immediately. In case of contact with skin, ninse immediately with cold running water for at least 15 minutes.

HOW TO STORE YOUR FIREROSE 5000 NIC SALTS

Store below 25 degrees Celsius in a clean dry place. Keep the product out of the sight and reach of children and animals.

ELUX Europe Ltd

Unit H9, Swan Business Centre, 4 Higher Swan Lane, Bolton BL3 3AQ, +44 (0) 1 204 773195

Regus Santry

The Crescent Building, Northwood Business Park, Santry, Dublin 9, Ireland, D09 C6X8, +353 1 893 4195

